



## Open Latarjet for instability

### **JARED M. MAHYLIS, MD**

Specialty Physicians of Illinois  
Olympia Fields, IL

#### **Phase I: (0 to 6 weeks after surgery)**

Goals: Protect Repair. Prevent dislocation, ensure wound healing.

1. Patients may remove dressing and shower 5 days after surgery. You may get the incision wet.
2. Sutures will dissolve on their own. No suture will need to be removed.
3. Ice is recommended to reduce swelling and help with pain. You should continue this for the first week at a minimum. Apply the ice to the surgical site 3-4 times per day once the nerve block has worn off.
4. You should maintain your sling when not performing exercises.
  - a. Initiate the following exercise program 3 times per day:
    - i. Immediate elbow, forearm, wrist, and hand range of motion out of sling . You may perform nonresistance, non-weight bearing use of the hand for activities such as typing on a computer or using a tablet computer.
    - ii. May dangle the arm to apply clothing and clean axilla.
    - iii. NO PASSIVE OR ACTIVE SHOULDER MOTION

#### **Phase II: (6 to 12 weeks after surgery)**

Goals: Initiate ROM & Protect Repair

1. You may continue sling use in public but should remove when comfortable and to begin shoulder motion .
2. No lifting weights or bearing weight on you arm.
3. Begin active assisted and passive ROM
  - a. start passive and active assistive ER at the side to 30 degrees and forward flexion to 120 degrees.
  - b. You may start active scapular mobility exercises at 4 weeks from surgery – Must keep the shoulder musculature relaxed.
  - c. No range of motion into abduction, ER >30 degrees or active IR

#### **Phase III: (3-6 months after surgery)**

Goal: Progress ROM and initiate strengthening.

1. May begin lifting up to 10 lbs below chest height and close to body and 5 lbs above chest height. Slow progression with goal to remove restrictions at 6 months .
2. Begin performing combined abduction and ER ROM and capsular mobility
3. Advance active and passive ROM in all planes to tolerance.
  - a. ER at the side and flexion to tolerance
  - b. Scapular plane elevation to Tolerance
  - c. IR and extension to tolerance
4. Initiate gentle rotator cuff strengthening.
5. Initiate scapular stabilizer strengthening



Jared M. Mahylis, MD

Jared M. Mahylis, MD  
Shoulder & Elbow Surgeon

**Phase IV: (>6 months after surgery)**

1. Stop all lifting restrictions.
2. Begin performing combined abduction and ER ROM and capsular mobility
3. Progress on your rotator cuff and scapular stabilizer strengthening.
4. Begin functional progression to sports specific activities
5. Continued functional strengthening